# CHLAA



**Test Booklet Code** 

GG

This Booklet contains 24 pages.

Do not open this Test Booklet until you are asked to do so.

Read carefully the Instructions on the Back Cover of this Test Booklet.

### Important Instructions:

- The Answer Sheet is inside this Test Booklet. When you are directed to open the Test Booklet, take out the Answer Sheet and fill in the particulars on Side-1 and Side-2 carefully with blue/black ball point pen only.
- The test is of 3 hours duration and this Test Booklet contains 180 questions. Each question carries 4 marks. For each correct response, the candidate will get 4 marks. For each incorrect response, one mark will be deducted from the total scores. The maximum marks are 720.
- 3. Use Blue/Black Ball Point Pen only for writing particulars on this page/marking responses.
- 4. Rough work is to be done on the space provided for this purpose in the Test Booklet only.
- On completion of the test, the candidate must hand over the Answer Sheet to the Invigilator before leaving the Room/Hall. The candidates are allowed to take away this Test Booklet with them.
- 6. The CODE for this Booklet is GG. Make sure that the CODE printed on Side-2 of the Answer Sheet is the same as that on this Test Booklet. In case of discrepancy, the candidate should immediately report the matter to the Invigilator for replacement of both the Test Booklet and the Answer Sheet.
- 7. The candidates should ensure that the Answer Sheet is not folded. Do not make any stray marks on the Answer Sheet. Do not write your Roll No. anywhere else except in the specified space in the Test Booklet/Answer Sheet.
- Use of white fluid for correction is not permissible on the Answer Sheet.

Offsets are produced by 8. The correct order of steps in Polymerase Chain Parthenocarpy Reaction (PCR) is (1)(2) Parthenogenesis Denaturation, Extension, Annealing Mitotic divisions (3) Denaturation, Annealing, Extension (4) Meiotic divisions Annealing, Extension, Denaturation (3)The experimental proof for semiconservative 2. Extension, Denaturation, Annealing replication of DNA was first shown in a India, the organisation responsible for (1) Plant assessing the safety of introducing genetically (2) Virus modified organisms for public use is (3)Bacterium (4) Fungus Research Committee Manipulation (RCGM) 3. Select the correct match : Genetic Engineering Appraisal Committee .(2) Matthew Meselson Pisum sativum (GEAC) and F. Stahl Council for Scientific and (3)Industrial (2)Francois Jacob and Lac operon Research (CSIR) Jacques Monod Indian Council of Medical Research (ICMR) Alfred Hershey and TMV Martha Chase Which of the following is commonly used as a Alec Jeffreys Streptococcus vector for introducing a DNA fragment in human pneumoniae lymphocytes? Which of the following pairs is wrongly à phage 4. matched? pBR 322 (1) XO type sex : Grasshopper (3) Ti plasmid Determination 1(4) Retrovirus (2)T.H. Morgan Linkage 11. Use of bioresources by multinational companies (3) ABO blood grouping : Co-dominance and organisations without authorisation from the Starch synthesis in pea (4) : Multiple alleles concerned country and its people is called 5. Select the correct statement: Biodegradation Spliceosomes take part in translation. . (2)r Bioexploitation (2)Transduction was discovered by S. Altman. (3)Biopiracy (3) Punnett square was developed by a British Bio-infringement (4)scientist. A 'new' variety of rice was patented by a foreign Franklin Stahl coined the term "linkage". (4) company, though such varieties have been Which of the following has proved helpful in 6. present in India for a long time. This is related to preserving pollen as fossils? Lerma Rojo Oil content Basmati (2) Sporopollenin (3)Sharbati Sonora (3) Cellulosic intine (4)Co-667 (4) Pollenkitt

life-time?

Mango

Papaya

Jackfruit

Bamboo species

(I)

(2)

(3)

(4)

7.

Which of the following flowers only once in its

13.

(2)

Select the correct match:

F<sub>2</sub> × Recessive parent

T.H. Morgan

G. Mendel

Ribozymi

Transduction

Transformation

Dihybrid cross

Nucleic acid

Genetic

14.	Which of the following is true for nucleolus?	22.	Which one is wrongly matched?					
	(1) It takes part in spindle formation.		(1)	Gen	ıma cup	15	- Marchantia	
	(2) It is a site for active ribosomal RN	A	(2)	Uni	cellular	organ	nism - Chlorella	
- 10	synthesis.		(3)	Biffi	agellate	zoosp	oores - Brown algae	
1	(3) It is a membrane-bound structure.		(4)	Uni	flagellat	te gan	netes - Polysiphonia	
	(4) Larger nucleoli are present in dividing cell	S. 000	35.4		7	3	in Column Luish there in	
15.	The Golgi complex participates in	23.				2011/10/20	in Column I with those in	
	(1) Respiration in bacteria			Column II and select the correct option given below:				
	(2) Activation of amino acid		nen	Colu	me I		Column II	
	(3) Formation of secretory vesicles					-		
111	(4) Fatty acid breakdown	- 1	a.	Her	barium	1.	It is a place having a	
16.	Which of the following is not a product of light	ht					collection of preserved plants and animals.	
	reaction of photosynthesis?			17		11	A list that enumerates	
	(1) NADPH		b.	Key		ii.	methodically all the	
	(2) Oxygen						species found in an area	
ш	(3) NADH						with brief description	
	(4) ATP						aiding identification.	
17.	Which among the following is not a prokaryote	?	c.	Mus	eum	iii.	Is a place where dried and	
H	(1) Nostoc			212.000			pressed plant specimens	
	(2) Oscillatoria						mounted on sheets are	
	(3) Mycobacterium						kept.	
**	(4) Saccharomyces		d.	Cata	alogue	iv.	A booklet containing a list	
18.	Stomatal movement is <b>not</b> affected by						of characters and their	
	(1) O <sub>2</sub> concentration						alternates which are	
	(2) CO <sub>2</sub> concentration						helpful in identification of	
	(3) Light						various taxa.	
100	(4) Temperature			а	b	c	d	
19.		of	(1)	ii	iv	iii	1	
	sugars are		(2)	iii	iv	i	ii	
	(1) carbonyl and phosphate		(3)	iii	ii	i	iv	
	(2) carbonyl and hydroxyl		(4)	i	iv	iii	ii	
	(3) carbonyl and methyl		****	0000	**	earn or o		
(tax)	(4) hydroxyl and methyl	24.	Winged pollen grains are present in (1) Mango					
20.	The stage during which separation of the pairs homologous chromosomes begins is	ed	(1)	Pin				
	(1) Diakinesis		(3)	Cyc				
	(2) Zygotene		(4)	1	stard			
	(3) Diplotene	. 1	2000	278.000	1048.1 68			
	(4) Pachytene	25.	Afte	er kar	yogamy	follo	wed by meiosis, spores are	
21.	Stomata in grass leaf are	1	pro	duced	exogen	ously	in	
	(1) Rectangular		(1)	Aga	ricus			
	(2) Barrel shaped		(2)		charomy	vces		
	(3) Kidney shaped		(3)		rnaria			
	(4) Dumb-bell shaped		(4)	Neu	rospora			

Oxygen is not produced during photosynthesis by 33. 26. Cycas (1) the range of temperature that the organism Chara (2)needs to live (3) Nostne (2) the functional role played by the organism where it lives Green sulphur bacteria the physical space where an organism lives Double fertilization is all the biological factors in the organism's Fusion of two male gametes with one egg environment Syngamy and triple fusion Fusion of one male gamete with two polar 34. Which of the following is a secondary pollutant? nuclei SO. Fusion of two male gametes of a pollen tube (2) $O_2$ with two different eggs Which of the following elements is responsible for (3)COo 28. maintaining turger in cells? CO (4)Potassium Natality refers to (2)Calcium (1) Number of individuals leaving the habitat (3) Sodium Number of individuals entering a habitat (4) Magnesium (3) Birth rate 29. Which one of the following plants shows a very Death rate (4) close relationship with a species of moth, where none of the two can complete its life cycle without 36. World Ozone Day is celebrated on the other? (1) Banana 16th September (1)(2)Viola 22<sup>nd</sup> April (3) Yucca Hydrilla 21st April Pollen grains can be stored for several years in 5<sup>th</sup> June liquid nitrogen having a temperature of (1) - 196°C What type of ecological pyramid would obtained with the following data? -160°C (2)(3) -80°C Secondary consumer: 120 g (4) - 120°C Primary consumer: 60 g Primary producer: 10 g What is the role of NAD Upright pyramid of numbers (1) respiration? (1) It is a nucleotide source for ATP synthesis. Upright pyramid of biomass It is the final electron acceptor for anaerobic Pyramid of energy respiration. (4) Inverted pyramid of biomass It functions as an electron carrier. In stratosphere, which of the following elements It functions as an enzyme. 38. acts as a catalyst in degradation of ozone and 32. In which of the following forms is iron absorbed release of molecular oxygen? by plants? Fe (1) (1) Free element Oxygen Both ferric and ferrous (2)CI (3)Ferrous Carbon (4) Ferric

	(3)	Pericycle Epidermis			Colu	mnI		Column II	
40.	Plan	ts having little or no secondary growth are		a.	Glye	osuria ,	i	Accumulation of u	ric
1	(1)	Conifers Cycads		b.	Gou	1/	ii.	Mass of crystallise salts within the ki	
	(3) (4)	Deciduous angiosperms Grasses		c.	Ren	al calcul	1 /11		- 1
41.	(1) (2)	Carnivorous plants Submerged hydrophytes		d.		nerulap nritis	i¥.	arms out of Carrier Cours	e in
Н	(3) (4)	Free-floating hydrophytes Halophytes		(1)	a ii	b iii	e i	đ iv	
42.		et potato is a modified		190	iv	-4	ii	101	
201	(1)	Tap root		Van	10	-		Page 1	
ш	(2)	Rhizome		(3)	1	ii	iii	iv	
	(3)	Adventitious root		(4)	iii	ii	iv	1	
т	(4)	Stem	47.	Mat	ch the	items	given i	in Column I with th	hose in
43.	Whi	ch of the following statements is correct?		Column II and select the correct option gi					
	(1)	Horsetails are gymnosperms.		belo	W:				
П	(2)	Stems are usually unbranched in both Cycas and Cedrus.				imn I		Column II (Part of Exerc	etory
Н	(3)	Selaginella is heterosporous, while Salvinia is homosporous.						System)	
	(4)	Ovules are not enclosed by ovary wall in		n.	Ultr	afiltrati	on \	Henle's loop	
		gymnosperms.		b.	0.15.000	centrati	on	ii. Ureter	
44.		ct the wrong statement:			of u	rine	1	1	
1	(1)	Pseudopodia are locomotory and feeding structures in Sporozoans.		c.	Tran	nsport o e	f /	iii. Urinary blad	der
1	(2)	Mitochondria are the powerhouse of the cell in all kingdoms except Monera.		d.	Stor	age of u	rine	viv. Malpighian corpuscle	
	(3)	Mushrooms belong to Basidiomycetes.							
	(4)	Cell wall is present in members of Fungi and Plantae.						v. Proximal convoluted to	abule
45.		ndary xylem and phloem in dicot stem are luced by			a	b	e	d	
	(1)	Phellogen		(1)	V	iv	1	ii	
	(2)	Axillary meristems		(2)	v	īv	î	iii	
	(3)	Vascular cambium		1(3)	iv	i	ii	iii	
	(4)	Apical meristems		(4)	iv	v	ii	iii	

46.

below:

39. Casparian strips occur in

Cortex

Endodermis

(1)

(2)

Match the items given in Column I with those in

Column II and select the correct option given

	(3) Ascariasis (4) Elephantiasis	12	(3) (4)	Reptilia Amphibia
	(2) Amoebiasis		(2)	Osteichthyes
	(1) Ringworm disease	,	(1)	Aves
53.	In which disease does mosquito transmitted pathogen cause chronic inflammation of lymphatic vessels?		char	acterized by crop and gizzard in its digestive em.
	(4) Hemology	59.	Idon	tify the vertebrate group of animals
	(3) Analogy		(4)	using flagella for locomotion
	(2) Adaptive radiation		600	excess water
	(1) Convergent evolution		(3)	having a contractile vacuole for removing
32.	of many vertebrates is an example of		(2)	having two types of nuclei
52.	The similarity of bone structure in the forelimbs		(1)	using pseudopodia for capturing prey
	(4) Vitamin D	58.	Cilia	ates differ from all other protozoans in
	(3) Vitamin A		(4)	Dinoflagellates
	(2) Vitamin E	23	(3)	Diatoms
	(1) Vitamin B <sub>12</sub>		(2)	Euglenoids
51,	Conversion of milk to curd improves its nutritional value by increasing the amount of		(1)	Cyanobacteria
0	(4) b, c and e	53%	chie	f producers in the oceans?
	(3) a, b and c	57.	Whie	ch of the following organisms are known as
	(2) a, c and e			9 <sup>th</sup> abdominal segment
	(1) b, d and e		(4)	Presence of a boat shaped sternum on the
	e. Polygenic inheritance		(3)	Presence of caudal styles
	d. Incomplete dominance		(2)	Presence of anal cerci
	c. Multiple allele		(1)	Forewings with darker tegmina
	b. Co-dominance	00.		ale cockroach from a female cockroach?
	a. Dominance	56.	Whi	ch of the following features is used to identify
50.	Which of the following characteristics represent 'Inheritance of blood groups' in humans?		(4)	Macropus
	(4) Psoriasia	3	(3)	Chelone
-	(3) Rheumatoid arthritis		(2)	Psittacula
0	(2) Vitiligo		(1)	Camelus
	(1) Alzheimer's disease			neotherm?
49.	Which of the following is not an autoimmune disease?	55.	Whi	ch one of these animals is not a
	(4) Forelimbs of man, bat and cheetah		1(4)	Earthworm
	(3) Heart of bat, man and cheetah		(3)	Tunicate -
me	(2) Eye of octopus, bat and man		(2)	Starfish
	(1) Brain of bat, man and cheetah		(1)	Moth
20,	divergent evolution, select the incorrect option:	94.		ch of the following animals does not undergo amorphosis?
48.	Among the following sets of examples for	54	Whi	ch of the following animals does not undergo

60,	Hormones secreted by the placenta to maintain	64.	In a growing population of a country,					
	pregnancy are		(1) reproductive and pre-reproductive					
	(1) hCG, hPL, progestogens, estrogens		individuals are equal in number.  (2) pre-reproductive individuals are less than					
	(2) hCG, progestogens, estrogens,		the reproductive individuals.					
	glucocorticoids		(3) reproductive individuals are less than the					
	(3) hCG, hPL, estrogens, relaxin, oxytocin		post-reproductive individuals.					
	(4) hCG, hPL, progestogens, prolactin		(4) pre-reproductive individuals are more than the reproductive individuals.					
61.	The contraceptive 'SAHELI'	65.	Which part of poppy plant is used to obtain the drug "Smack" ?					
	(1) is an IUD.	- 69	(1) Roots					
Ш	(2) is a post-coital contraceptive.		(2) Leaves					
П	(3) increases the concentration of estrogen and		(3) Latex					
Ш	prevents ovulation in females.		(4) Flowers					
1	(4) blocks estrogen receptors in the uterus, preventing eggs from getting implanted.	66.	Match the items given in Column I with those in Column II and select the <i>correct</i> option given below:					
62.	The amnion of mammalian embryo is derived		Column II Column II					
	from		a. Eutrophication i. UV-B radiation					
	(1) mesoderm and trophoblast		b. Sanitary landfill ii. Deforestation					
	(2) ectoderm and endoderm		c. Snow blindness iii. Nutrient					
	(3) endoderm and mesoderm		enrichment					
Н	(4) ectoderm and mesoderm		d. Jhum cultivation iv. Waste disposal a b c d					
Ш			(1) iii iv i ii					
63.	The difference between spermiogenesis and		(2) i ii iv iii					
	spermiation is		(3) i iii iv ii					
	(1) In spermiogenesis spermatozoa from sertoli		(4) ii i iii iv					
	cells are released into the cavity of seminiferous tubules, while in spermitation spermatozoa are formed.	67.	Which one of the following population interactions is widely used in medical science for the production of antibiotics?					
	(2) In spermiogenesis spermatozoa are formed,		(1) Parasitism					
	while in spermiation spermatozoa are		(2) Amensalism					
	released from sertoli cells into the cavity of		(3) Mutualism (4) Commensalism					
	seminiferous tubules.		Acti Settimativa (MATZO)					
	(3) In spermiogenesis spermatozoa are formed, while in spermiation spermatids are	15.5500	All of the following are included in 'Ex-situ conservation' except					
	formed.		(1) Botanical gardens					
	(4) In spermiogenesis spermatids are formed,		(2) Seed banks					
	while in spermiation spermatozoa are		(3) Sacred groves					
	formed.		(4) Wildlife safari parks					

69.	Which of the following gastric cells indirectly help in erythropoiesis?				73.	73. Which of the following is an amino acid derived hormone?					
	(1)	Goblet cells				(1)	Estradiol				
1	(2)	Parietal cells				(2)	Estriol				
	(3)	Mucous cells				(3)	Ecdysone				
	(4)	Chief cells			١.,	Var	Epinephrine				
70.	Mate	h the items o	riven in	Column I with those in	1	(CSE)	Ершеригию				
1,725		mn II and se		e correct option given	74.		ch of the following		uctures or regions is function?		
		Column I		Column II		(1)	Hypothalamus	:	production of		
	a.	Fibrinogen	10	Osmotic balance					releasing hormones and regulation of		
	b.	Globulin	TT:	Blood clotting					temperature, hunger and thirst.		
	c.	Albumin	Hi.	Defence mechanism		(2)	Corpus callosum		band of fibers		
	(1)	a b i iii	e ii			(4)	Corpus canasum		connecting left and right cerebral hemispheres.		
	(2)	ii iii	i			vor	Timber materi	0401	CC 15000 - 1510 1000 100		
	(3)	i ii	iii		3	101	Limbic system	-20	consists of fibre tracts that		
	(4)	iii ii	i						interconnect different regions of		
71.	Calci	ium is imp	portant se it	in skeletal muscle					brain; controls movement.		
	(1)			head from the actin		(4)	Medulla oblongata	:	controls respiration and cardiovascular reflexes.		
	(2)			tion of bonds between oridges and the actin	75.		ch of the followin		ormones can play a sis?		
	(3)		myosin	ATPase by binding to	1	(I)	Estrogen and Para	athy	roid hormone		
		it					(2) Parathyroid hormone and Prolactin				
	(4)	(4) binds to troponin to remove the masking of active sites on actin for myosin.				(3)	Progesterone and				
1						(4)	Aldosterone and Prolactin				
72.		th of the firatory disord		r is an occupational	76.		transparent lens in dace by	the	human eye is held in		
	(1) Botulism					(1)	smooth muscles at	tach	ed to the iris		
	(2)	Emphysema				(2)	(2) smooth muscles attached to the ciliary				
	(3)	Silicosis				(3)	ligaments attache	d to	the iris		
	(4)	Anthracis			7.	(4)	ligaments attache	d to	the ciliary body		

Nissl bodies are mainly composed of Which of the following options correctly 77. represents the lung conditions in asthma and Nucleic acids and SER emphysema, respectively? (2) Free ribosomes and RER Increased respiratory surface; (3) DNA and RNA Inflammation of bronchioles Proteins and lipids Decreased respiratory surface; Inflammation of bronchioles Which of these statements is incorrect? 81. Increased number of bronchioles; Increased Glycolysis operates as long as it is supplied respiratory surface with NAD that can pick up hydrogen atoms. Inflammation of bronchioles; Decreased Oxidative phosphorylation takes place in respiratory surface outer mitochondrial membrane. Match the items given in Column I with those in Glycolysis occurs in cytosol. (8) Column II and select the correct option given Enzymes of TCA cycle are present in (4) below: mitochondrial matrix. Column II Column I Many ribosomes may associate with a single Between left atrium Tricuspid valve a. mRNA to form multiple copies of a polypeptide and left ventricle simultaneously. Such strings of ribosomes are Between right b. Bicuspid valve termed as wentricle and Plastidome (1)pulmonary artery Nucleosome (2)Semilunar valve iii. Between right atrium and right Polyhedral bodies (3) ventricle Polysome Which of the following terms describe human 83. iii dentition? (2)iii Pleurodont, Monophyodont, Homodont (1) (3)iii Pleurodont, Diphyodont, Heterodont (2)17 (3) Thecodont, Diphyodont, Heterodont Match the items given in Column I with those in Thecodont, Diphyodont, Homodont Column II and select the correct option given 84. Which of the following events does not occur in below: rough endoplasmic reticulum? Column II Column I Cleavage of signal peptide Tidal volume 2500 - 3000 mL Phospholipid synthesis Inspiratory Reserve 1100 - 1200 mL (3)Protein glycosylation volume Protein folding (4) Expiratory Reserve Hi. 500 - 550 mL volume 85. Select the incorrect match: Residual volume iv. 1000 - 1100 mL Submetacentric - L-shaped chromososmes chromosomes Occytes of amphibians Polytene (2)(1) ii iii chromosomes (2)Шi Sex chromosomes (3)Allosomes (3) iii iv ii. Diplotene bivalents Lampbrush (4) chromosomes (4) iii i iv SPACE FOR ROUGH WORK CHLAA/GG/Page 9 English

- 86. AGGTATCGCAT is a sequence from the coding strand of a gene. What will be the corresponding sequence of the transcribed mRNA?

  (1) ACCUAUGCGAU

  (2) UCCAUAGCGUA

  (3) UGGTUTCGCAT

  (4) AGGUAUCGCAU
- According to Hugo de Vries, the mechanism of evolution is
  - (1) Phenotypic variations
  - (2) Minor mutations
  - (3) Saltation
  - (4) Multiple step mutations
- 88. Match the items given in Column I with those in Column II and select the correct option given below:

Column I Column II Proliferative Phase i. Breakdown of endometrial lining Secretory Phase Follicular Phase Menstruation Luteal Phase iii (2)ii (3)iii ii

- 89. All of the following are part of an operon except
  - (1) an enhancer

iii

(4)

- (2) a promoter
- (3) structural genes
- (4) an operator
- A woman has an X-linked condition on one of her X chromosomes. This chromosome can be inherited by
  - (1) Only grandchildren
  - (2) Both sons and daughters
  - (3) Only sons
  - (4) Only daughters


- 91. A tuning fork is used to produce resonance in a glass tube. The length of the air column in this tube can be adjusted by a variable piston. At room temperature of 27°C two successive resonances are produced at 20 cm and 73 cm of column length. If the frequency of the tuning fork is 320 Hz, the velocity of sound in air at 27°C is
  - (1) 350 m/s
  - (2) 300 m/s
  - (3) 339 m/s
  - (4) 330 m/s
- 92. The electrostatic force between the metal plates of an isolated parallel plate capacitor C having à charge Q and area A, is
  - proportional to the square root of the distance between the plates.
  - (2) inversely proportional to the distance between the plates.
  - (3) linearly proportional to the distance between the plates.
  - (4) independent of the distance between the plates.
- 93. An electron falls from rest through a vertical distance h in a uniform and vertically upward directed electric field E. The direction of electric field is now reversed, keeping its magnitude the same. A proton is allowed to fall from rest in it through the same vertical distance h. The time of fall of the electron, in comparison to the time of fall of the proton is
  - (1) 10 times greater
  - (2) equal
  - (3) 5 times greater
  - (4) smaller
- 94. A pendulum is hung from the roof of a sufficiently high building and is moving freely to and fro like a simple harmonic oscillator. The acceleration of the bob of the pendulum is 20 m/s<sup>2</sup> at a distance of 5 m from the mean position. The time period of oscillation is
  - (1) 2s
  - (2) 1s
  - (3) ms
  - (4) 2π s

- 95. Current sensitivity of a moving coil galvanometer is 5 div/mA and its voltage sensitivity (angular deflection per unit voltage applied) is 20 div/V. The resistance of the galvanometer is
  - (1) 250 Ω
  - (2) 500 Ω
  - (3) 25 Ω
  - (4) 40 Ω
- 96. A metallic rod of mass per unit length 0.5 kg m<sup>-1</sup> is lying horizontally on a smooth inclined plane which makes an angle of 30° with the horizontal. The rod is not allowed to slide down by flowing a current through it when a magnetic field of induction 0.25 T is acting on it in the vertical direction. The current flowing in the rod to keep it stationary is
  - (1) 14·76 A
  - (2) 11·32 A
  - (3) 5.98 A
  - (4) 7-14 A
- 97. A thin diamagnetic rod is placed vertically between the poles of an electromagnet. When the current in the electromagnet is switched on, then the diamagnetic rod is pushed up, out of the horizontal magnetic field, Hence the rod gains gravitational potential energy. The work required to do this comes from
  - the lattice structure of the material of the rod
  - (2) the induced electric field due to the changing magnetic field
  - (3) the magnetic field
  - (4) the current source
- 98. An inductor 20 mH, a capacitor 100 μF and a resistor 50 Ω are connected in series across a source of emf, V = 10 sin 314 t. The power loss in the circuit is
  - (1) 2.74 W
  - (2) 1·13 W
  - (3) 0.43 W
  - (4) 0.79 W


- 99. An object is placed at a distance of 40 cm from a concave mirror of focal length 15 cm. If the object is displaced through a distance of 20 cm towards the mirror, the displacement of the image will be
  - 1) 30 cm towards the mirror
  - (2) 36 cm towards the mirror
  - (3) 36 cm away from the mirror
  - (4) 30 cm away from the mirror
- 100. An em wave is propagating in a medium with a velocity V = Vî. The instantaneous oscillating electric field of this em wave is along +y axis. Then the direction of oscillating magnetic field of the em wave will be along
  - (1) y direction
  - (2) x direction
  - (3) + z direction
  - (4) z direction
- A thin diamagnetic rod is placed vertically between the poles of an electromagnet. When the current in the electromagnet is switched on, then inductor is 60 mA. This inductor is of inductance
  - (1) 1-389 H
  - (2) 13-89 H
  - (3) 138·88 H
  - (4) 0·138 H
  - 102. The refractive index of the material of a prism is √2 and the angle of the prism is 30°. One of the two refracting surfaces of the prism is made a mirror inwards, by silver coating. A beam of monochromatic light entering the prism from the other face will retrace its path (after reflection from the silvered surface) if its angle of incidence on the prism is
    - (1) 30°
    - (2) zero
    - (3) 45°
    - (4) 60°

- 103. The ratio of kinetic energy to the total energy of 107. In the circuit shown in the figure, the input an electron in a Bohr orbit of the hydrogen atom,
  - (1) 2:-1
  - (2)1:-2
  - 1:-1 (3)
  - (4) 1:1
- 104. An electron of mass m with an initial velocity  $\overrightarrow{V} = \overrightarrow{V_0} \hat{i} (V_0 > 0)$  enters an electric field  $E = -E_0 \hat{i}$  ( $E_0 = constant > 0$ ) at t = 0. If  $\lambda_0$  is its de-Broglie wavelength initially, then its de-Broglie wavelength at time t is
  - (1) \(\lambda\_0 \) t
  - (2) \(\lambda\_0\)
  - (3)  $\lambda_0 \left[ 1 + \frac{eE_0}{mV_0} t \right]$
  - $(4) \frac{\lambda_0}{\left(1 + \frac{eE_0}{mV_0}t\right)}$
- 105. For a radioactive material, half-life 10 minutes. If initially there are 600 number of nuclei, the time taken (in minutes) for the disintegration of 450 nuclei is
  - (1) 30
  - (2)15
  - (3)10
  - (4) 20
- 106. When the light of frequency  $2v_0$  (where  $v_0$  is threshold frequency), is incident on a metal plate, the maximum velocity of electrons emitted is v1. When the frequency of the incident radiation is increased to 5vo, the maximum velocity of electrons emitted from the same plate is v2. The ratio of v1 to v2 is
  - (1) 4:1
  - 2:1 (2)
  - (3)1:4
  - (4) 1:2

voltage  $V_i$  is 20 V,  $V_{HE} = 0$  and  $V_{CE} = 0$ . The values of I<sub>B</sub>, I<sub>C</sub> and β are given by




- (1)  $I_B = 20 \mu A$ ,  $I_C = 5 mA$ ,  $\beta = 250$
- (2)  $I_B = 40 \mu A$ ,  $I_C = 5 mA$ ,  $\beta = 125$
- (3)  $I_B = 25 \mu A$ ,  $I_C = 5 m A$ ,  $\beta = 200$
- (4)  $I_B = 40 \mu A$ ,  $I_C = 10 mA$ ,  $\beta = 250$
- 108. In a p-n junction diode, change in temperature due to heating
  - (1) does not affect resistance of p-n junction
  - (2) affects the overall V I characteristics of p-n junction
  - (3)affects only forward resistance
  - affects only reverse resistance
- 109. In the combination of the following gates the output Y can be written in terms of inputs A and Bas





- A.B + A.B
- A + B
- $A, \overline{B} + \overline{A}, B$
- (4)

- rings of different colours for identification. The colour code sequence will be
  - Yellow Green Violet Gold
  - Green Orange Violet Gold
  - Yellow Violet Orange Silver
  - Violet Yellow Orange Silver
- 111. A set of 'n' equal resistors, of value 'R' each, are connected in series to a battery of emf 'E' and internal resistance 'R'. The current drawn is I. Now, the 'n' resistors are connected in parallel to the same battery. Then the current drawn from battery becomes 10 I. The value of 'n' is
  - 20 (1)
  - (2)
  - (3)11
  - (4) 10
- 112. A battery consists of a variable number 'n' of identical cells (having internal resistance 'r' each) which are connected in series. The terminals of the battery are short-circuited and the current I is measured. Which of the graphs shows the correct relationship between I and n?










- 110. A carbon resistor of (47 ± 4·7) kΩ is to be marked | 113. The efficiency of an ideal heat engine working between the freezing point and boiling point of water, is
  - 6.25% (1)
  - 12-5% (2)
  - (3)20%
  - 26-8% (4)
  - 114. The volume (V) of a monatomic gas varies with its temperature (T), as shown in the graph. The ratio of work done by the gas, to the heat absorbed by it, when it undergoes a change from state A to state B, is



- (1)
- (2)
- (3)
- (4)
- 115. The fundamental frequency in an open organ pipe is equal to the third harmonic of a closed organ pipe. If the length of the closed organ pipe is 20 cm, the length of the open organ pipe is
  - 12-5 cm (1)
  - (2)16 cm
  - (3) 8 cm
  - (4) 13.2 cm
- 116. At what temperature will the rms speed of oxygen molecules become just sufficient for escaping from the Earth's atmosphere? (Given:

Mass of oxygen molecule (m) =  $2.76 \times 10^{-26}$  kg Boltzmann's constant  $k_B = 1.38 \times 10^{-23} \text{ J K}^{-1}$ 

- $5.016 \times 10^4 \text{ K}$
- $1.254 \times 10^4 \text{ K}$
- $8.360 \times 10^4 \text{ K}$
- $2.508 \times 10^4 \text{ K}$


- 117. The power radiated by a black body is P and it radiates maximum energy at wavelength,  $\lambda_0$ . If the temperature of the black body is now changed so that it radiates maximum energy at wavelength  $\frac{3}{4}\lambda_0$ , the power radiated by it becomes nP. The value of n is
  - (1)  $\frac{256}{81}$
  - (2)  $\frac{81}{256}$
  - (3)  $\frac{4}{3}$
  - $(4) \frac{3}{4}$
- 118. Two wires are made of the same material and have the same volume. The first wire has cross-sectional area A and the second wire has cross-sectional area 3A. If the length of the first wire is increased by Δl on applying a force F, how much force is needed to stretch the second wire by the same amount?
  - (1) 4F
  - (2) F
  - (3) 6 F
  - (4) 9 F
- 119. A small sphere of radius 'r' falls from rest in a viscous liquid. As a result, heat is produced due to viscous force. The rate of production of heat when the sphere attains its terminal velocity, is proportional to
  - (1) r<sup>5</sup>
  - (2) r<sup>4</sup>
  - (3) r<sup>2</sup>
  - (4) r<sup>3</sup>
- 120. A sample of 0·1 g of water at 100°C and normal pressure (1·013 × 10<sup>5</sup> Nm<sup>-2</sup>) requires 54 cal of heat energy to convert to steam at 100°C. If the volume of the steam produced is 167-1 cc, the change in internal energy of the sample, is
  - (1) 42·2 J
  - (2) 84·5 J
  - (3) 208·7 J
  - (4) 104·3 J

121. The kinetic energies of a planet in an elliptical orbit about the Sun, at positions A, B and C are K<sub>A</sub>, K<sub>B</sub> and K<sub>C</sub>, respectively. AC is the major axis and SB is perpendicular to AC at the position of the Sun S as shown in the figure. Then



- (1) K<sub>R</sub> < K<sub>A</sub> < K<sub>C</sub>
- (2) K<sub>B</sub> > K<sub>A</sub> > K<sub>C</sub>
- (3) K<sub>A</sub> > K<sub>B</sub> > K<sub>C</sub>
- (4) K<sub>A</sub> < K<sub>B</sub> < K<sub>C</sub>
- 122. A solid sphere is in rolling motion. In rolling motion a body possesses translational kinetic energy (K<sub>t</sub>) as well as rotational kinetic energy (K<sub>r</sub>) simultaneously. The ratio K<sub>t</sub>: (K<sub>t</sub> + K<sub>r</sub>) for the sphere is
  - (1) 10:7
  - (2) 2:5
  - (3) 5:7
  - (4) 7:10
- 123. A solid sphere is rotating freely about its symmetry axis in free space. The radius of the sphere is increased keeping its mass same. Which of the following physical quantities would remain constant for the sphere?
  - Rotational kinetic energy
  - (2) Angular momentum
  - (3) Moment of inertia
  - (4) Angular velocity
- 124. If the mass of the Sun were ten times smaller and the universal gravitational constant were ten times larger in magnitude, which of the following is not correct?
  - Time period of a simple pendulum on the Earth would decrease.
  - (2) 'g' on the Earth will not change.
  - Walking on the ground would become more difficult.
  - (4) Raindrops will fall faster.


125. A body initially at rest and sliding along a frictionless track from a height h (as shown in the figure) just completes a vertical circle of diameter AB = D. The height h is equal to



- (1)  $\frac{7}{5}$  D
- (2)  $\frac{5}{4}$  D
- (3) D
- (4)  $\frac{3}{2}$  D
- 126. Three objects, A: (a solid sphere), B: (a thin circular disk) and C: (a circular ring), each have the same mass M and radius R. They all spin with the same angular speed ω about their own symmetry axes. The amounts of work (W) required to bring them to rest, would satisfy the relation
  - (1)  $W_B > W_A > W_C$
  - (2) W<sub>A</sub> > W<sub>C</sub> > W<sub>B</sub>
  - (3)  $W_A > W_B > W_C$
  - (4)  $W_C > W_B > W_A$
- 127. A moving block having mass m, collides with another stationary block having mass 4m. The lighter block comes to rest after collision. When the initial velocity of the lighter block is v, then the value of coefficient of restitution (e) will be
  - (1) 0-8
  - (2) 0.4
  - (3) 0.25
  - (4) 0.5
- 128. Which one of the following statements is 132.

  incorrect?
  - Frictional force opposes the relative motion.
  - (2) Coefficient of sliding friction has dimensions of length.
  - (3) Limiting value of static friction is directly proportional to normal reaction.
  - (4) Rolling friction is smaller than sliding friction.

- 129. The moment of the force,  $\overrightarrow{F} = 4 \hat{i} + 5 \hat{j} 6 \hat{k}$  at (2, 0, -3), about the point (2, -2, -2), is given by
  - (1)  $-7\hat{i} 8\hat{j} 4\hat{k}$
  - (2) -7î-4ĵ-8k
  - (3)  $-4\hat{i} \hat{j} 8\hat{k}$
  - (4)  $-8\hat{i} 4\hat{j} 7\hat{k}$
  - 130. A toy car with charge q moves on a frictionless horizontal plane surface under the influence of a uniform electric field E. Due to the force q E, its velocity increases from 0 to 6 m/s in one second duration. At that instant the direction of the field is reversed. The car continues to move for two more seconds under the influence of this field. The average velocity and the average speed of the toy car between 0 to 3 seconds are respectively.
    - (1) 1 m/s, 3·5 m/s
    - (2) 1-5 m/s, 3 m/s
    - (3) 1 m/s, 3 m/s
    - (4) 2 m/s, 4 m/s
- 131. A block of mass m is placed on a smooth inclined wedge ABC of inclination θ as shown in the figure. The wedge is given an acceleration 'a' towards the right. The relation between a and θ for the block to remain stationary on the wedge is



- (1)  $a = g \cos \theta$
- (2)  $a = g \tan \theta$
- (3)  $a = \frac{g}{\sin \theta}$
- (4)  $a = \frac{g}{\cos \alpha c \theta}$
- 132. A student measured the diameter of a small steel ball using a screw gauge of least count 0-001 cm. The main scale reading is 5 mm and zero of circular scale division coincides with 25 divisions above the reference level. If screw gauge has a zero error of - 0-004 cm, the correct diameter of the ball is
  - (1) 0.053 cm
  - (2) 0.529 cm
  - (3) 0.525 cm
  - (4) 0.521 cm

- 133. Unpolarised light is incident from air on a plane surface of a material of refractive index 'μ'. At a particular angle of incidence 'i', it is found that the reflected and refracted rays are perpendicular to each other. Which of the following options is correct for this situation?
  - (1)  $i = \sin^{-1}\left(\frac{1}{\mu}\right)$
  - (2)  $i = tan^{-1} \left(\frac{1}{\mu}\right)$
  - (3) Reflected light is polarised with its electric vector perpendicular to the plane of incidence
  - (4) Reflected light is polarised with its electric vector parallel to the plane of incidence
- 134. In Young's double slit experiment the separation d between the slits is 2 mm, the wavelength λ of the light used is 5896 Å and distance D between the screen and slits is 100 cm. It is found that the angular width of the fringes is 0·20°. To increase the fringe angular width to 0·21° (with same λ and D) the separation between the slits needs to be changed to
  - (1) 2·1 mm
  - (2) 1-7 mm
  - (3) 1-9 mm
  - (4) 1.8 mm
- 135. An astronomical refracting telescope will have large angular magnification and high angular resolution, when it has an objective lens of
  - (1) large focal length and large diameter
  - (2) small focal length and small diameter
  - (3) large focal length and small diameter
  - (4) small focal length and large diameter

- 136. In which case is the number of molecules of water maximum?
  - 0.00224 L of water vapours at 1 atm and 273 K
  - (2) 10<sup>-3</sup> mol of water
  - (3) 0.18 g of water
  - (4) 18 mL of water
- 137. Consider the change in oxidation state of Bromine corresponding to different emf values as shown in the diagram below:

$$BrO_4^- \xrightarrow{1.82 \text{ V}} BrO_3^- \xrightarrow{1.5 \text{ V}} HBrO$$

$$Br^- \xleftarrow{1.0652 \text{ V}} Br_2 \xleftarrow{1.595 \text{ V}}$$

Then the species undergoing disproportionation is

- (1) Br<sub>2</sub>
- (2) HBrO
- (3) BrO,
- (4) BrO.
- Among CaH<sub>2</sub>, BeH<sub>2</sub>, BaH<sub>2</sub>, the order of ionic character is
  - (1) BeH<sub>2</sub> < BaH<sub>2</sub> < CaH<sub>2</sub>
  - (2) BaH<sub>2</sub> < BeH<sub>2</sub> < CaH<sub>2</sub>
  - (3) CaH<sub>2</sub> < BeH<sub>2</sub> < BaH<sub>2</sub>
  - (4) BeH<sub>2</sub> < CaH<sub>2</sub> < BaH<sub>2</sub>
- 139. The correct difference between first- and second-order reactions is that
  - a first-order reaction can be catalyzed; a second-order reaction cannot be catalyzed
  - (2) the rate of a first-order reaction does depend on reactant concentrations; the rate of a second-order reaction does not depend on reactant concentrations
  - (3) the half-life of a first-order reaction does not depend on [A]<sub>0</sub>; the half-life of a second-order reaction does depend on [A]<sub>0</sub>
  - (4) the rate of a first-order reaction does not depend on reactant concentrations; the rate of a second-order reaction does depend on reactant concentrations

4.49	-					H WO	RK English
(4)	iv	v	ii	i		(4)	B < Al < In < Ga < Tl
	1		iii			(3)	B < Al < Ga < In < Tl
(2)	iii	v	i	ii		(2)	B < Ga < Al < In < Tl
(1)	iv	1	ii	iii		(1)	B < Ga < Al < Tl < In
	a	b	c v.	d d.m.	150.		correct order of atomic radii in group 13 nents is
Ma	141					(4)	Fe
						(3)	Zn
c.			iii.	$\sqrt{3}$ B.M.	1	(2)	Cu
b.	Cr <sup>3+</sup>		ii.	$\sqrt{35}$ B.M.		(1)	Mg
a,	$\mathrm{Co}^{3+}$		i.	√8 B.M.	1000000		owing metals can be used to reduce alumina?
Column I				Column II	149	Con	sidering Ellingham diagram, which of the
spin magnetic moments of the ions given in Column II and assign the correct code:						(4)	All form monobasic oxyacids.
네 드리마 방송 아르는데 아들아 내일 때 아름이 내려왔다면서 아이를 잃었다. 그리 말이 없는데 아이는 이는 아이는데 그리고 있다고 있다며 아이를 다 했다.						(3)	All are oxidizing agents.
(4)	(4) tetranuclear					(2)	Chlorine has the highest electron-gain enthalpy.
(3)	2) dinuclear					(3.55)	states.
(2)						(1)	All but fluorine show positive exidation
							ich of the following statements is <b>not</b> true for ogens?
						1,40	22103, 210, 213, 211401
							HNO <sub>3</sub> , NO, N <sub>2</sub> , NH <sub>4</sub> Cl
							HNO <sub>3</sub> , NO, NH <sub>4</sub> Cl, N <sub>2</sub>
(1)		75-200				(2)	NH <sub>4</sub> Cl, N <sub>2</sub> , NO, HNO <sub>3</sub>
			The state of the s			(1)	HNO <sub>3</sub> , NH <sub>4</sub> Cl, NO, N <sub>2</sub>
The	geome	etry a	nd magn	etic behaviour of the			correct order of N-compounds in its reasing order of oxidation states is
(4)	CrO <sub>4</sub>						one
(3)	Cr <sub>2</sub> O	7					two
						(2)	three
						(1)	four
			CO TOTAL CONTRACTOR	•	2.400		lectrons on central atom 'Cl' is
							he structure of CIF3, the number of lone pairs
						(4)	Ga
(3) Coordination isomerism						200	Al
(2)		11.					In
(1)	Ioniz	ation is	somerism			1200	В
The type of isomerism shown by the complex [CoCl <sub>2</sub> (en) <sub>2</sub> ] is							n MF <sub>6</sub> <sup>3-</sup> ion?
	(2) (3) (4) Which d-d (1) (2) (3) (4) The conn (1) (2) (3) (4) Iron (1) (2) (3) (4) Match spin Column Column column d. (1) (2) (3) (4) (4) (1) (2) (3) (4) (4) (1) (2) (3) (4) (4) (1) (2) (3) (4) (1) (2) (3) (4) (1) (2) (3)	(2) Links (3) Coord (4) Geom Which on d-d transit (1) MnO (2) MnO (3) Cr <sub>2</sub> O (4) CrO <sub>4</sub> The geome complex [N (1) squar (2) tetral (3) tetral (4) squar Iron carbor (1) trinuc (2) dinuc (3) mono (4) tetral Match the spin magn Column II	(2) Linkage ison (3) Coordinatio (4) Geometrical Which one of d-d transition and (1) MnO <sub>4</sub> <sup>-</sup> (2) MnO <sub>4</sub> <sup>2-</sup> (3) Cr <sub>2</sub> O <sub>7</sub> <sup>2-</sup> (4) CrO <sub>4</sub> <sup>2-</sup> The geometry at complex [Ni(CO) <sub>4</sub> (1) square plant (2) tetrahedral (3) tetrahedral (4) square plant (2) trinuclear (3) mononuclear (3) mononuclear (4) tetranuclear (3) mononuclear (4) tetranuclear (5) mononuclear (6) tetranuclear (7) mononuclear (8) mononuclear (9) dinuclear (1) trinuclear (1) trinuclear (2) dinuclear (3) mononuclear (4) tetranuclear (5) mononuclear (6) tetranuclear (7) mononuclear (8) mononuclear (9) dinuclear (1) trinuclear (1) trinuclear (2) dinuclear (3) mononuclear (4) tetranuclear (5) mononuclear (6) tetranuclear (7) dinuclear (8) mononuclear (9) dinuclear (1) trinuclear (1) trinuclear (2) dinuclear (3) mononuclear (4) tetranuclear (5) dinuclear (6) dinuclear (7) dinuclear (8) mononuclear (9) dinuclear (1) trinuclear (1) trinuclear (2) dinuclear (3) mononuclear (4) tetranuclear (5) dinuclear (6) dinuclear (7) dinuclear (8) dinuclear (9) dinuclear (1) trinuclear (1) trinuclear (2) dinuclear (3) mononuclear (4) tetranuclear (6) dinuclear (7) dinuclear (8) dinuclear (9) dinuclear (1) trinuclear (1) trinuclear (2) dinuclear (3) mononuclear (4) tetranuclear (6) dinuclear (7) dinuclear (8) dinuclear (9) dinuclear (1) trinuclear (1) trinuclear (2) dinuclear (3) mononuclear (4) tetranuclear (5) dinuclear (6) dinuclear (7) dinuclear (8) dinuclear (9) dinuclear (1) trinuclear (1) trinuclear (1) trinuclear (2) dinuclear (3) dinuclear (4) tetranuclear (6) dinuclear (7) dinuclear (8) dinuclear (9) dinuclear (1) dinuclear (1) trinuclear (2) dinuclear (3) dinuclear (4) tetranuclear (6) dinuclear (7) dinuclear (8) dinuclear (9) dinuclear (1) dinuclear (1) dinuclear (2) dinuclear (3) dinuclear (4) tetranuclear	(2) Linkage isomerism (3) Coordination isomeris (4) Geometrical isomeris Which one of the fold-distransition and parama (1) MnO <sub>4</sub> (2) MnO <sub>4</sub> (3) Cr <sub>2</sub> O <sub>7</sub> (4) CrO <sub>4</sub> (4) CrO <sub>4</sub> (5) The geometry and magnitude complex [Ni(CO) <sub>4</sub> ] are (1) square planar geome (2) tetrahedral geometry (3) tetrahedral geometry (4) square planar geome (5) trinuclear (6) trinuclear (7) dinuclear (8) mononuclear (9) dinuclear (10) magnetic moments (11) Column I  12) a. Co <sup>3+</sup> 13. b. Cr <sup>3+</sup> 14. c. Fe <sup>3+</sup> 15. c. Fe <sup>3+</sup> 16. Ni <sup>2+</sup> 17. c. Fe <sup>3+</sup> 18. d. Ni <sup>2+</sup> 19. d. Ni <sup>2</sup>	(2) Linkage isomerism (3) Coordination isomerism (4) Geometrical isomerism Which one of the following ions exhibits d-d transition and paramagnetism as well? (1) MnO₄ (2) MnO₄ (3) Cr₂O₂ (4) CrO₄ (4) CrO₄ (5) Square planar geometry and paramagnetic geometry and paramagnetic geometry and paramagnetic geometry and diamagnetic geometry geometry and diamagnetic geometry geometry and diamagnetic geometry geometry and diamagnetic geometry geom	(2) Linkage isomerism (3) Coordination isomerism (4) Geometrical isomerism (4) Geometrical isomerism Which one of the following ions exhibits d-d transition and paramagnetism as well? (1) MnO₄ (2) MnO₄ (3) Cr₂O₂ (4) CrO₄ (4) CrO₄ (5) The geometry and magnetic behaviour of the complex [Ni(CO)₄] are (1) square planar geometry and paramagnetic (2) tetrahedral geometry and diamagnetic (3) tetrahedral geometry and diamagnetic (4) square planar geometry and diamagnetic (4) square planar geometry and diamagnetic Iron carbonyl, Fe(CO)₅ is (1) trinuclear (2) dinuclear (3) mononuclear (4) tetranuclear  Match the metal ions given in Column I with the spin magnetic moments of the ions given in Column II and assign the correct code:  Column I Column II 149.  a. Co³+ i. √8 B.M. b. Cr³+ ii. √35 B.M. c. Fe³+ iii. √35 B.M. d. Ni²+ iv. √24 B.M. v. √15 B.M. a b c d (1) iv i ii iii (2) iii v i ii iii (3) i iii iii iii	(1) Ionization isomerism (2) Linkage isomerism (3) Coordination isomerism (4) Geometrical isomerism (4) Geometrical isomerism (5) Minch one of the following ions exhibits d-d transition and paramagnetism as well? (1) MnO₄ (2) MnO₄ (2) MnO₄ (3) Cr₂O₂ (4) CrO₄ (4) CrO₄ (5) Cr₂O₂ (6) Cro₄ (6) Square planar geometry and paramagnetic (1) square planar geometry and paramagnetic (2) tetrahedral geometry and diamagnetic (3) tetrahedral geometry and diamagnetic (4) square planar geometry and diamagnetic (3) tetrahedral geometry and diamagnetic (4) square planar geometry and diamagnetic (3) trinuclear (4) trinuclear (5) dinuclear (6) trinuclear (7) dinuclear (8) mononuclear (9) dinuclear (1) trinuclear (1) trinuclear (2) dinuclear (3) mononuclear (4) tetranuclear (4) tetranuclear (5) Match the metal ions given in Column I with the spin magnetic moments of the ions given in Column II and assign the correct code:  Column I Column II  a. Co₃⁴ i. √8 B.M. b. Cr³⁺ ii. √35 B.M. c. Fe³⁺ iii. √35 B.M. c. Fe³⁺ iii. √3 B.M. d. Ni²⁺ iv. √24 B.M. v. √15 B.M. a b c d (1) iv i ii iii (2) iii v i ii (3) (3) (4) (4) (5) The elect (1) (7) The column II (2) (8) iii iii iii (9) iii v i ii (10) (2) (11) (2) (2) (12) (2) (33) (3) (4)

#### 151. In the reaction

the electrophile involved is

- (1) dichloromethyl anion (CHCl<sub>2</sub>)
- (2) dichlorocarbene (:CCl2)
- (3) formyl cation (CHO)
- (4) dichloromethyl cation (CHCl<sub>2</sub>)

### 152. Carboxylic acids have higher boiling points than aldehydes, ketones and even alcohols of comparable molecular mass. It is due to their

- more extensive association of carboxylic acid via van der Waals force of attraction
- (2) formation of intermolecular H-bonding
- (3) formation of carboxylate ion
- (4) formation of intramolecular H-bonding


## 153. Compound A, C<sub>8</sub>H<sub>10</sub>O, is found to react with NaOI (produced by reacting Y with NaOH) and yields a yellow precipitate with characteristic smell.

A and Y are respectively

(1) 
$$CH - CH_3$$
 and  $I_2$   
OH

- 154. Which of the following molecules represents the order of hybridisation sp<sup>2</sup>, sp<sup>2</sup>, sp, sp from left to right atoms?
  - (1) CH2 = CH CH = CH2
  - (2)  $CH_3 CH = CH CH_3$
  - (3) CH<sub>2</sub> = CH C = CH
  - (4) HC = C C = CH

### 155. Which of the following carbocations is expected to be most stable?



- 156. Which of the following is correct with respect to -I effect of the substituents ? (R = alkyl)
  - (1) -NH<sub>2</sub> > OR > F
  - (2) -NR<sub>2</sub>>-OR>-F
  - (3) −NR<sub>2</sub> < −OR < −F</p>
  - (4) -NH2 <-OR <-F

- 157. Regarding cross-linked or network polymers, which of the following statements is incorrect?
  - Examples are bakelite and melamine.
  - They contain strong covalent bonds in their polymer chains.
  - They are formed from bi- and tri-functional (3)monomers.
  - (4) They contain covalent bonds between various linear polymer chains.
- 158. Nitration of aniline in strong acidic medium also gives m-nitroaniline because
  - always goes to m-position.
  - In acidic (strong) medium aniline is present as anilinium ion.
  - In electrophilic substitution (3)reactions amino group is meta directive.
  - In spite of substituents nitro group always goes to only m-position.
- 159. The difference between amylose and amylopectin
  - (1) Amylopectin have  $1 \rightarrow 4$   $\alpha$ -linkage and 1 → 6 β-linkage
  - (2) Amylose is made up of glucose and galactose
  - Amylose have  $1 \rightarrow 4$   $\alpha$ -linkage 1 → 6 β-linkage
  - (4) Amylopectin have 1 -> 4 α-linkage and  $1 \rightarrow 6 \alpha$ -linkage
- 160. A mixture of 2.3 g formic acid and 4.5 g oxalic acid is treated with conc. HoSO4. The evolved gaseous mixture is passed through KOH pellets. Weight (in g) of the remaining product at STP will be
  - (1) 2.8
  - (2) 4.4
  - (3) 3.0
  - 1.4
- 161. Which of the following exides is most acidic in nature?
  - (1) BaO
  - (2) CaO
  - (3) BeO
  - (4) MgO

162. For the redox reaction

$$MnO_4^- + C_2O_4^{2-} + H^+ \longrightarrow Mn^{2+} + CO_2 + H_2O$$

the correct coefficients of the reactants for the balanced equation are

$MnO_4^-$	$C_2O_4^{2-}$	$H^{+}$
	-30	915

- (1)
- 2 (2)5 16
- (3)2 16 (4) 16
- (1) In absence of substituents nitro group 163. Which one of the following conditions will favour maximum formation of the product in the reaction.

$$A_2(g) + B_2(g) \rightleftharpoons X_2(g)$$
  $\Delta_r H = -X \text{ kJ } ?$ 

- (1) High temperature and high pressure
- High temperature and low pressure
- (3) Low temperature and low pressure
- (4) Low temperature and high pressure
- 164. When initial concentration of the reactant is doubled, the half-life period of a zero order reaction
  - (1) is tripled
  - remains unchanged (2)
  - is doubled (3)
  - is halved
- 165. The bond dissociation energies of X<sub>2</sub>, Y<sub>2</sub> and XY are in the ratio of 1:0.5:1. AH for the formation of XY is -200 kJ mol-1. The bond dissociation energy of X2 will be
  - (1) 800 kJ mol-1
  - (2) 400 kJ mol-1
  - 100 kJ mol-1 (3)
  - 200 kJ mol-1
- 166. The correction factor 'a' to the ideal gas equation corresponds to
  - electric field present between the gas molecules
  - forces of attraction between the (2) molecules
  - volume of the gas molecules
  - (4) density of the gas molecules

- 167. Following solutions were prepared by mixing different volumes of NaOH and HCl of different concentrations:
  - a.  $60 \text{ mL } \frac{\text{M}}{10} \text{ HCl} + 40 \text{ mL } \frac{\text{M}}{10} \text{ NaOH}$
  - b.  $55 \text{ mL } \frac{M}{10} \text{ HCl} + 45 \text{ mL } \frac{M}{10} \text{ NaOH}$
  - c.  $75 \text{ mL } \frac{\text{M}}{5} \text{ HCl} + 25 \text{ mL } \frac{\text{M}}{5} \text{ NaOH}$
  - d. 100 mL  $\frac{M}{10}$  HCl + 100 mL  $\frac{M}{10}$  NaOH

pH of which one of them will be equal to 1?

- (1) d
- (2) c
- (3) a
- (4) b
- 168. On which of the following properties does the coagulating power of an ion depend?
  - Both magnitude and sign of the charge on the ion
  - (2) The sign of charge on the ion alone
  - (3) Size of the ion alone
  - (4) The magnitude of the charge on the ion alone
- 169. Given van der Waals constant for NH<sub>3</sub>, H<sub>2</sub>, O<sub>2</sub> and CO<sub>2</sub> are respectively 4·17, 0·244, 1·36 and 3·59, which one of the following gases is most easily liquefied?
  - (1) O<sub>2</sub>
  - (2) CO<sub>9</sub>
  - (3) H<sub>2</sub>
  - (4) NH<sub>3</sub>
- 170. The solubility of  ${\rm BaSO_4}$  in water is  $2\cdot42\times10^{-3}~{\rm gL^{-1}}$  at 298 K. The value of its solubility product ( ${\rm K_{sp}}$ ) will be

(Given moIar mass of  $BaSO_4 = 233 \text{ g mol}^{-1}$ )

- (1)  $1.08 \times 10^{-14} \text{ mol}^2 \text{ L}^{-2}$
- (2)  $1.08 \times 10^{-8} \text{ mol}^2 \text{ L}^{-2}$
- (3)  $1.08 \times 10^{-12} \text{ mol}^2 \text{ L}^{-2}$
- (4)  $1.08 \times 10^{-10} \text{ mol}^2 \text{ L}^{-2}$

171. Identify the major products P, Q and R in the following sequence of reactions:

$$\begin{array}{c} & + \operatorname{CH_3CH_2CH_2Cl} \xrightarrow{Ah \text{ydrous}} \\ & \xrightarrow{AlCl_3} \\ & \\ P \xrightarrow{(i) \ O_2} \\ & \xrightarrow{(ii) \ H_3O^+/\Delta} Q + R \end{array}$$

$$(2) \quad \bigcirc \stackrel{CH(CH_3)_2}{\longleftarrow} \quad \bigcirc \stackrel{OH}{\longleftarrow} \quad CH_3 - CO - CH_3$$

(3) 
$$CH_2CH_2CH_3$$
 CHO COOH

(4) 
$$CH_2CH_2CH_3$$
  $CHO$   
,  $CH_3CH_2-OH$ 

- 172. Which of the following compounds can form a zwitterion?
  - (1) Benzoic acid
  - (2) Glycine
  - (3) Acetanilide
  - (4) Aniline

173. The compound C7H8 undergoes the following 177. Consider the following species: reactions:

$$C_7H_8 \xrightarrow{3 Cl_2/\Delta} A \xrightarrow{Br_2/Fe} B \xrightarrow{Zn/HCl} C$$

The product 'C' is

- (1) 3-bromo-2,4,6-trichlorotoluene
- (2) p-bromotoluene
- o-bromotoluene (3)
- (4) m-bromotoluene
- 174. Which exide of nitrogen is not a common pollutant introduced into the atmosphere both due to natural and human activity?
  - N<sub>9</sub>O
  - (2) NO
  - (3) NO<sub>0</sub>
  - (4) NoOn
- 175. The compound A on treatment with Na gives B, and with PCI5 gives C. B and C react together to give diethyl ether. A, B and C are in the order
  - C<sub>9</sub>H<sub>8</sub>Cl, C<sub>9</sub>H<sub>6</sub>, C<sub>9</sub>H<sub>5</sub>OH
  - (2) C<sub>2</sub>H<sub>5</sub>OH, C<sub>2</sub>H<sub>5</sub>ONa, C<sub>2</sub>H<sub>5</sub>Cl
  - (3) C<sub>2</sub>H<sub>5</sub>OH, C<sub>2</sub>H<sub>5</sub>Cl, C<sub>2</sub>H<sub>5</sub>ONa
  - (4) C<sub>2</sub>H<sub>5</sub>OH, C<sub>2</sub>H<sub>6</sub>, C<sub>2</sub>H<sub>5</sub>Cl
- 176. Hydrocarbon (A) reacts with bromine by substitution to form an alkyl bromide which by Wurtz reaction is converted to gaseous hydrocarbon containing less than four carbon atoms, (A) is
  - (1) CH<sub>3</sub> CH<sub>3</sub>
  - (2) CH<sub>4</sub>
  - (3) CH<sub>2</sub> = CH<sub>2</sub>
  - (4) CH = CH

CN+, CN-, NO and CN

Which one of these will have the highest bond order?

- (1) CN<sup>†</sup>
- (2) CN
- CN
- (4) NO
- 178. Magnesium reacts with an element (X) to form an ionic compound. If the ground state electronic configuration of (X) is 1s2 2s2 2p3, the simplest formula for this compound is
  - (1) MgoX
  - (2) Mg<sub>3</sub>X<sub>2</sub>
  - (3) MgX<sub>2</sub>
  - Mg<sub>2</sub>X<sub>2</sub>
- 179. Iron exhibits bcc structure at room temperature. Above 900°C, it transforms to fcc structure. The ratio of density of iron at room temperature to that at 900°C (assuming molar mass and atomic radii of iron remains constant with temperature)
  - (1)
  - (2)
  - (3)
  - (4)
- Which one is a wrong statement?
  - (1) The electronic configuration of N atom is

$$\begin{array}{cccc} 1s^2 & 2s^2 & 2p_x^1 & 2p_y^1 & 2p_z^1 \\ \uparrow \downarrow & \uparrow \downarrow & \uparrow \uparrow \downarrow & \\ \end{array}$$

- (2) The value of m for d<sub>2</sub> is zero.
  - An orbital is designated by three quantum numbers while an electron in an atom is designated by four quantum numbers.
  - Total orbital angular momentum of electron in 's' orbital is equal to zero.